

JOURNAL OF THE MEDICAL WOMEN'S ASSOCIATION OF NIGERIA

Established in 2004

March 2025

Volume 10: No 1

Original Article

Access this article online

Quick Response Code:



Website: www.jmwan.org.ng

DOI: <https://doi.org/10.71526/jmwan.v10i1.66>

¹Department of Paediatrics
Faculty of Clinical sciences,
University of Nigeria Enugu
Campus
Enugu state

²Department of Paediatrics
University of Nigeria Teaching
Hospital, Ituku Ozalla
Enugu state.

³Department of Paediatrics
Nnamdi Azikiwe University Awka

⁴Department of Paediatrics
Nnamdi Azikiwe University
Teaching Hospital Nnewi

⁵Department of Orthopaedics
University of Nigeria Teaching
Hospital, Ituku Ozalla
Enugu state.

Address for correspondence:

Odimegwu CL
Department of Paediatrics
Faculty of Clinical sciences,
University of Nigeria Enugu
Campus, Enugu state
chioma.odimegwu@unn.edu.ng
+2347033030952

Comparative Assessment of Neutrophil Counts and use of Neutropenia to predict Low CD4 Count in Anti-Retroviral Therapy (Art)-Naive HIV positive Children in Enugu, Nigeria

Chime PU^{1,2}, Odimegwu CL^{1,2}, Onyia JOT², Nwaneli E^{3,4}, Okoro KA⁵

Abstract

BACKGROUND: Neutropenia has been noted to be associated with HIV disease progression and reduced survival in children. Neutropenia has been postulated to be associated with low CD4 count in HIV-positive children by some researchers. Hence, this study sought to assess the prevalence of neutropenia in ART-naïve HIV-positive patients and compare them to HIV-negative subjects and also to investigate if reduced absolute neutrophil counts can be used to predict low CD4 count in the HIV-positive patients.

METHODS: This study was a prospective observational study carried out at the University of Nigeria Teaching Hospital, Enugu, involving a total of 200 children (100 test subjects and 100 test controls). Biodata was collected with a proforma and blood samples were subsequently collected and analysed for neutrophil count and CD₄ count. Data generated were analysed with Statistical Package for Social Sciences (SPSS) version 22.

RESULTS: The result showed that neutropenia with a prevalence of 18% is not significantly associated with HIV infections. It was also shown that there is no significant association between the age of the children and neutropenia. Likewise, CD4 count was not significantly associated with neutropenia.

CONCLUSION: The study showed that neutropenia in the subjects is comparable with that of the controls and cannot be used as a substitute for CD₄ count or as one of the criteria for commencement of antiretroviral medications in ART-naïve HIV-positive children.

Keywords:

HIV, antiretroviral therapy naïve, neutropenia, CD₄ count, children.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Introduction

The Human Immunodeficiency Virus (HIV) is one of the most important emerging infections of this century.^[1] The HIV infection causes the Acquired Immunodeficiency Syndrome (AIDS), which is a systemic disorder characterised by deficiency in cellular immune responses.^[1] It is probably one of the diseases with multiple impacts on persons, families, communities and the entire society.^[1] HIV is threatening, especially in sub-Saharan African countries. In Nigeria, the prevalence rose from 3.8% in 2003 to 5.8% in 2011.^[1] Haematological complications such as neutropenia have been documented to be the second most common cause of morbidity and mortality in HIV-positive persons.^[2] A variety of haematological manifestations are seen at every stage of HIV and often life-threatening and impair the quality of life of these patients.^[3]

Haematological abnormalities such as neutropenia have been documented as strong independent predictors of morbidity and mortality in HIV-infected children.^[3] Although it is not part of the criteria for initiating therapy nor used by the World Health Organisation (WHO) for staging HIV, peripheral blood cell abnormalities in an abnormal haemogram are important prognostic tools for morbidity in HIV infection and AIDS.^[3] The 2011 Mbanya *et al.*^[4] study at the Central Hospital of Yaoundé, done among ART-naïve AIDS patients, revealed that 95.4% of AIDS patients had morphological blood cell disorders compared to 32.2% amongst healthy controls. Haematological manifestations in HIV-infected patients often pose a great challenge in the comprehensive management of such patients and may cause symptoms that are life-threatening and impair the quality of life of these patients.^[5]

The haematological complications, such as neutropenia, are associated with HIV disease progression and subsequent reduced survival.^[5] It has been noted that a large decline in neutrophil count in HIV-positive patients is a pointer for later development of HIV-induced central nervous system (CNS) disease, which is by all means catastrophic to the patients.^[6] This complication in HIV-infected ART-naïve patients

usually results in poor ART treatment outcomes when they eventually become eligible for ART and strongly predicts poor prognosis and increased mortality.^[6] This complication, when present, must be addressed appropriately and in good time.

Neutropenia is defined as a neutrophil count less than 1500 cells/mm³ and is the most common leucopenia occurring in HIV-infected individuals.^[7] Neutropenia is common and its incidence rises from 13% to 44% with disease progression from HIV to AIDS.^[7] HIV infection suppresses the bone marrow and leads to decreased levels of granulocyte colony-stimulating factor, the factor that stimulates production of white blood cells in the bone marrow and affects the granulocyte-macrophage lineage, resulting in leucopenia and neutropenia.^[8] Neutropenia can also result from other opportunistic infections, including cytomegalovirus, tuberculosis, histoplasmosis and leishmaniasis.^[15] Hence, HIV patients with cytopenias may require bone marrow examination to determine the cause and to direct therapy.^[8]

Low CD₄ count is defined as a CD₄ count less than 500 cells/mm³. Low CD₄ count has been associated with prevalent anaemia, neutropenia and thrombocytopenia.^[9] HIV preferentially infects CD₄ cells, causing their destruction.^[10] It has been calculated that each day, more than 1 billion CD₄ cells are destroyed.^[10] The decline in CD₄ count is linked to viral load and is used as a measure of disease progression.^[10]

Neutropenia has been associated with a low baseline CD₄ count.^[11] Taking into account the patient's CD₄ cell count, Suresh *et al.*^[11] showed that the incidence of neutropenia was 0.8% in HIV patients with CD₄ count greater than 700 cells/mm³ and rose to 13.4% in those with CD₄ count less than 250 cells/mm³.

Hence, it is concluded that there is a strong negative association between CD₄ counts and the severity of neutropenia, anaemia and thrombocytopenia in HIV patients.^[12] Therefore, a complete blood count with differential should be performed every three months to screen for haematological abnormalities in HIV-infected children. A complete blood count may need to be obtained more often for children receiving bone

marrow suppressive therapy or if abnormalities are identified.

Although neutropenia is a common manifestation of HIV infection and may have considerable impact on patients' well-being, treatment and care, few studies on neutrophil counts in HIV-infected children have been undertaken in Africa, but none in Nigeria in particular and Enugu in general, to the best of the author's knowledge. Such information for HIV-infected children in Enugu may help to forestall potential treatment failure later.

By way of justification, Africa is the epicentre of the HIV epidemic, as reported by the World Health Organisation (WHO), and Sub-Saharan Africa contributes about 67% of the burden worldwide.^[1] About 22 million people were living with the virus by the end of 2007.^[1] Nigeria has the largest number of people living with HIV/AIDS in West Africa and the second largest in the subcontinent.^[1] Neutropenia is associated with HIV disease progression and subsequent reduced survival.^[2] It has been noted that a large decline in neutrophil counts in HIV-positive patients is a pointer for later development of HIV-induced CNS disease, which is by all means catastrophic to the patients.^[6] This complication in HIV-infected ART-naïve patients usually results in poor ART treatment outcomes when they eventually become eligible for ART and strongly predicts poor prognosis and increased mortality. This complication, when present, must be addressed appropriately and in good time. Studies on neutrophil counts in HIV-infected children are almost non-existent in Nigeria, hence the compelling need to carry out this vital and essential study and hopefully go a long way in averting the potential problems this might cause to the patients. This study will also provide baseline information for the country as well as provide data on which further studies on the topic can be developed.

The aim of this research is to study the effects of human immunodeficiency virus (HIV) infection on neutrophil count in HIV-positive ART-naïve children seen at University of Nigeria Teaching Hospital (UNTH), Enugu, while the objectives are as follows: to assess the level of neutrophil counts in ART-naïve HIV-positive patients and compare it to the uninfected patients (control); to determine the prevalence of neutropenia in ART-naïve HIV-positive patients; and to relate the

level of neutrophil counts with the level of CD4 counts in ART-naïve HIV-positive patients.

Materials and Methods

Study Design: This was a prospective observational study conducted at a tertiary health facility with paediatric HIV/AIDS facilities: the University of Nigeria Teaching Hospital, Enugu. Paediatric HIV/AIDS clinics usually run once a week. Subjects were enrolled as they presented to the clinic for the first time or for a check-up. Information about biodata, medical data and demography were obtained from all the children under study through their parents or guardians using a proforma designed by the researcher. Subjects and controls were physically examined and their axillary temperature checked with a mercury thermometer to exclude those with febrile conditions. Their weights were also measured in kilograms and their weight-for-age centiles were used to determine their nutritional status and hence exclude those that were malnourished using modified welcome classification. Neutrophil counts were determined on subjects and controls.

Study Population:

Subjects: All ART-naïve HIV positive children

Controls: All HIV negative children

Sample Size Estimation

Sample selection was by consecutive enrolment.

Sample size was calculated using the formula^[15]:

$$n = \frac{Z^2 pq}{d^2}.$$

Where n = Sample Size

Z = 1.96 for 95% Confidence interval

P = Proportion with the characteristic (HIV positive subjects) = 5.8% (WHO, 2012).

q = Proportion without the characteristic = 1 – 0.058

= 0.942

d = Degree of accuracy, usually set at 0.05.

$$n = \frac{1.96^2 \times 0.058 \times 0.942}{0.05^2}$$

$$n = 83.96$$

Attrition rate = 10% of calculated sample size

Minimal sample size = 92.36

However, sample size of 100 was used.

Inclusion Criteria:

1. All HIV-positive ART-naïve children who were attending the Paediatric HIV/AIDS clinic at UNTH, Enugu.
2. All HIV-positive ART-naïve children of consenting mothers.

Exclusion Criteria:

1. HIV-positive children who have commenced ART.
2. HIV-positive ART-naïve children with inconsistent neutrophil counts.
3. Malnourished or febrile HIV-positive ART-naïve children.
4. HIV positive ART-naïve children on medications that have the potential to cause myelosuppression such as anti-cancer drugs.

Data Presentation and Analysis: Data generated were entered into the proforma by the researcher and were subsequently entered into a computer data base. The data were then analyzed with Statistical Package for Social Sciences (SPSS)

version 22 of the IBM Corporation, 1 New Orchard Rd, Armonk, NY 10504.

Descriptive statistics such as frequency, percentages, mean, median and standard deviations were used to summarise categorical and continuous variables. Median values of neutrophil count (skewed variable) were compared between subjects and age/sex-matched controls using the Mann-Whitney U Test. Relationships between age, CD4 count and neutrophil counts were done using correlation and linear regression. Associations between HIV status, different levels of CD4 count and neutrophil counts were tested using logistic regression and chi-square. The statistical significance was set at $p < 0.05$. Results were presented in tables and figures.

Laboratory Methods: After obtaining an informed consent from the mothers/guardians, blood samples (about 2 mL each) were collected from peripheral veins into an EDTA-containing sample bottle and rocked severally to allow for proper mixture of blood and anticoagulant. The samples were then analysed by the researcher using the autoanalyser under the supervision of a laboratory scientist. Another blood sample, 2 mL each, were collected two weeks after from the same subjects, and subjects with consistent values on both occasions were enrolled.

Ethical Consideration: The approval of the Health Research Ethics Committee of the University of Nigeria Teaching Hospital, Enugu was obtained. Informed consent to participate was obtained from all of the participants during investigations.

Results

Table 1: Sex and age distribution of subjects and controls

Age (group)	Subjects		Control	
	Male	Female	Male	Female
1 – 3	19 (39.6)	14 (26.9)	19 (39.6)	14 (26.9)
4 – 6	9 (18.8)	15 (28.8)	9 (18.8)	15 (28.8)
7 – 9	14 (29.2)	16 (30.8)	14 (29.2)	16 (30.8)
10 – 12	6 (12.5)	7 (13.5)	6 (12.5)	7 (13.5)
Total	48 (100)	52 (100)	48 (100)	52 (100)

$$\chi^2 = 2.332, p = 0.431$$

A total of 200 children (100 subjects and 100 controls) were enrolled into the study. All matched for sex and age. There were 48 males

and 52 females with ratio of approximately 1:1 but the gender difference was not statistically significant. ($p = 0.431$). The ages of the children

were grouped as shown in Table 1. The age ranged from 1 to 12 years with a mean $\pm$ standard deviation (SD) of 5.59 ± 3.25 years. The

predominant ages were in the 1-3 years group while the least predominant ages were in the 10-12 years group. (Table 1)

Table 2: Comparison of absolute neutrophil count in the Subjects and Controls

Parameters	Subjects Median (Mean rank)	Controls Median (Mean rank)	Mann-Whitney U	p-value
Neutrophil counts	2428.50 (90.51)	3320.50 (110.50)	4000.50	0.015

In the subjects, the Neutrophil Counts ranged from 572 to 18711 cells/mm³ with a mean $\pm$ standard deviation (SD) of 3176 ± 2469 cells/mm³. In the controls, the Neutrophil Counts ranged from 105 to 12450 cells/mm³ with a mean $\pm$ standard deviation of 3966.14 ± 2646.14 cells/mm³.

Therefore, a non-parametric alternative of t-test which is the Mann-Whitney U test was used to determine the association of ANC between the subjects and controls.

Median neutrophil count for subjects (2428.50) when compared to that of controls (3320.00) were significantly different. (p = 0.015). (Table 2)

The neutrophil counts were not normally distributed in both the subjects and the controls.

Table 3: Prevalence of neutropenia and its association with Subjects and Controls.

	Subject	Control	p value	OR	95% C.I for OR
<i>Neutrophil Counts</i>					
Neutropenia	18	14	0.441	1.348	0.630 – 2.887
Normal	82	86			

Neutropenia was seen in 18 subjects (11 males and 7 females) and in 14 controls (9 males and 5 females). Prevalence of neutropenia is 18% among the subjects and 14% among the controls.

There is no significant association in the prevalence of neutropenia between the subjects and the controls (p = 0.441). (Table 3)

Table 4: Comparison of absolute neutrophil count in Male and Female Subjects and Controls

Parameters	Gender		Mann-Whitney U	p-value
	Male Median (Mean rank)	Female Median (Mean rank)		
Neutrophil count (Subjects)	2394.00 (48.05)	2472.50 (52.76)	1130.50	0.418
Neutrophil count (Controls)	3138.00 (46.29)	(3407.00 (54.08)	1048.50	0.181

In the subjects, there is no significant difference between: Gender and Absolute Neutrophil Count, p = 0.418. In the controls also, there is no significant difference between: Gender and Absolute Neutrophil Count, p = 0.181. (Table 4)

Table 5: Relationship between age and neutrophil counts in subjects and controls

Age	Statistics	Neutrophil counts (cell/mm ³) Subjects	Neutrophil counts (cell/mm ³) Controls
	Pearson Correlation (r)	-0.110	0.046
	p value	0.277	0.651
	N	100	100

There is no significant relationship between the neutrophil counts and age in both the Subjects and the Controls. (Table 5)




Figure 1: Prevalence of different levels of CD₄ count in the subjects

The overall minimum CD₄ count was 4 cell/mm³ and the maximum was 2098 cells/mm³ with a median CD₄ count of 350 cells/mm³. Eighteen percent of the study population had CD₄ counts of <200 cells. Nineteen percent of them had CD₄ counts between 200 and 500 cells. Majority of the study population (63%) had CD₄ above 500 cells. (Figure 1)

Table 6: Relationship between levels of absolute neutrophil count and CD₄ count of the Subjects

Statistics	Neutrophil counts
R (Correlation Coefficient)	0.119
R ² (Coefficient of Determination)	0.014
B (Regression coefficient)	0.565
p value	0.239

There is no significant relationship between CD₄ count and absolute neutrophil count, $p = 0.239$.

Discussion

Infection with HIV ultimately leads to profound immunosuppression in which patients may present with various AIDS defining clinical conditions that may impact negatively on the quality of life of the patients. Impaired haematopoiesis, immune mediated cytopenias and altered coagulation mechanisms have all been described in HIV infected children. These abnormalities may occur as a result of HIV infection, as sequelae of HIV-related opportunistic infections or malignancies. Neutropenia and other haematological

manifestations have been documented to be the second most common cause of morbidity and mortality in HIV patients.^[9] A study was conducted to identify haematological parameters which could serve as pointers to the diagnosis of HIV/AIDS in resource poor settings in view of the fact that the cost of diagnostic and monitoring techniques for HIV/AIDS are unaffordable to most people in resource poor settings.^[6]

This prospective observational study assessed the haematological abnormalities such as neutropenia in HIV positive ART-naïve children (subjects) and compares with the HIV negative children (controls).

The prevalence of neutropenia in this study was 18% in the subjects and 14% in the controls. This 18% prevalence was higher in studies done by Erhabor *et al.*,^[12] and Amballie *et al.*,^[6] which showed prevalence of 6% and 10% respectively. The difference may be due to variation in study population, clinical conditions and study design methods.

In this study, more males had greater percentage of neutropenia than the females both in the subjects, 11(61%) Vs 7(39%) and in the controls, 9(64%) Vs 5(36%) but the differences were not statistically different.

The increase or decrease in ages of both the subjects and the controls had no effect on the neutrophil counts. This contrasted with the studies by Erhabor *et al.*^[12] and Amballie *et al.*^[6]

In this study, there was no significant relationship between neutropenia and CD₄ counts. This was in contrast to the studies by Calenda and Chermann,^[14] who reported a significant association between neutropenia and CD₄ counts.

Conclusion

This study shows that there is a significant difference in neutrophil count between the HIV-positive children and their negative counterparts. It was also noted that a lower CD₄ count could not be predicted by neutropenia. It was therefore recommended that a low neutrophil count cannot be used as one of the criteria for commencing ART, especially in resource-poor climes where estimation of CD₄ count is difficult.

Conflict of interest statement: The authors report no conflicts of interest.

Funding: This study was not funded by any organization. We bore all the expenses that accrued from in study.

Acknowledgments: We are grateful to the research assistant who helped in data entering.

References

1. Abrams DI, Steinhart C, Frascino R. Epoetin alfa therapy for anaemia in HIV-infected patients: impact on quality of life. *International journal of STD & AIDS*. 2000 Oct 1;11(10):659-65.
2. Addis Z, Yitayew G. Prevalence of some haematological abnormalities among HIV positive patients on their first visit to a tertiary health institution in Ethiopia; A cross sectional study. *Int Bld Res Rev*. 2014, 2: 270-278.
3. Adetifa, IMO., Teniye EO, Akinsulie AO, Ezeaka VC. Hematological abnormalities associated with pediatric HIV/AIDS in Lagos. *Ann Trop Pediatr: Int. Child Health*. 2016, 26:121-125.
4. Mbanya DNS, Tsafack JP, Binam F, Afane Ze E, Kaptue LN. **Peripheral blood cell changes in AIDS patients in Yaoundé, Cameroon.** *Bull Liais Doc OCEAC*. 2011, 34: 14-17.
5. Alcantara S. Thrombocytopenia is strongly associated with simian AIDS in pigtail macaques. *J Acquir Immune Defic Syndr*. 2019, 51: 374-379.
6. Amballie AA, Ajibola AA, Ogun SA, Ogunkolo OF. Demographic pattern and haematological profile in people living with HIV/AIDS in a university teaching hospital. *Scie Essay*. 2017, 2: 315-318.
7. Centre for Disease Control and Prevention DoHA. Revised classification system for human immunodeficiency virus infection in children under 13 years of age. *MMWR* 2014, 43:1–10.
8. Castaldo A, Tarallo L, Palomba E. Iron deficiency and intestinal malabsorption in HIV disease. *J Pediatr Gastroenterol Nutr*. 2016, 22: 359-363.
9. Cosby CD. Haematologic disorders associated with HIV and AIDS. *J Infus Nurs*. 2017, 30: 22-32.
10. Denué BA, Gashua W, Bello HS. Relation between some haematological abnormalities in treatment-naïve HIV infected patients. *East Medit Hlth J*. 2013, 19: 362-368.
11. Suresh VS, Singh VP, Madhuka R. Hematological profile of HIV patients in relation to immune status-a hospital based cohort from Varanasi, North India. *Turk J Hematol*. 2018, 25: 13-19.
12. Erhabor O, Ejele OA, Nwauche CA. Some haematological parameters in HIV infected Africans: the Nigerian perspective. *Niger J Med*. 2015, 14: 33-38.
13. Ferede, Wondimeneh. Prevalence and related factors of anaemia in HAART-naïve positive patients at Gondar University Hospital, Northwest Ethiopia. *BMC Hematology*. 2013, 13:8.
14. Calenda V, Chermann JC. The effects of HIV on haematopoiesis. *Euro J Haematol*. 2012, 48: 181-186.
15. Aroye MO. *Research Methodology with Statistics for Health and Social Sciences*. Ilorin: Nathadex Publishers. 2013, 115-129.